

Tooth eruption The primary teeth

We don't usually think of a newborn as having teeth. However, at birth the crowns of the 20 "baby" or primary teeth are almost completely formed, and they are hidden from view in an infant's jawbones. The primary teeth gradually erupt through the gums during the first 2½ years of life.

The four front teeth—two upper and two lower—usually erupt first, beginning as early as six months after birth. Most children have a full set of primary teeth by the time they are 3 years old. The child's jaws continue to grow, making room for the permanent (adult) teeth that will begin to erupt at about age 6 years. Primary teeth begin to shed between ages 6 and 7 years. This process continues until about age 12 years.

The chart and photograph identify the names of the primary teeth and provide the approximate ages at which you can expect the teeth to erupt and shed.

Primary teeth may be temporary, but they deserve good care. A child needs strong, healthy primary teeth not only to chew food easily, but to pronounce words properly.

This first set of teeth also holds a place in the jaw for the permanent teeth, which move into place as the primary teeth are shed. Primary teeth should be kept clean and healthy so that a child can remain free of cavities and oral pain. Infection from decayed primary teeth can damage the permanent teeth developing under them.

Parents and other caregivers may not realize that primary teeth are susceptible to decay as soon as they appear in the mouth. Tooth decay in infants and toddlers sometimes is called early childhood caries, baby bottle tooth decay or nursing mouth syndrome. This condition can destroy teeth. It occurs when a child's teeth are exposed frequently to sugary liquids for long periods.

You can help reduce the risk of tooth decay. Never allow your infant or toddler to fall asleep with a bottle containing milk, formula, fruit juices or sweetened liquid. Don't dip a pacifier in sugar or honey. If your infant or toddler needs a comforter between regular feedings or at bedtime, give the child a clean pacifier recommended by your dentist or pediatrician.

Primary Teeth			
	Upper Teeth	Erupt	Shed
	Central Incisor	8-12 Months	6-7 Years
	Lateral Incisor	9-13 Months	7-8 Years
	Canine (Cuspid)	16-22 Months	10-12 Years
	First Molar	13-19 Months	9-11 Years
	Second Molar	25-33 Months	10-12 Years
	Lower Teeth	Erupt	Shed
	Second Molar	23-31 Months	10-12 Years
	First Molar	14-18 Months	9-11 Years
	Canine (Cuspid)	17-23 Months	9-12 Years
	Lateral Incisor	10-16 Months	7-8 Years
	Central Incisor	6-10 Months	6-7 Years



Wipe your child's gums with a wet washcloth or a clean gauze pad after each feeding. Begin brushing your child's teeth with a little water as soon as the first tooth appears. Supervise toothbrushing to make sure that children older than 2 years use only a pea-sized amount of fluoride toothpaste and avoid swallowing it. Children should be taught to spit out remaining toothpaste and rinse with water after brushing. ■

Prepared by the ADA Division of Communications, in cooperation with The Journal of the American Dental Association and the ADA Council on Scientific Affairs. Unlike other portions of JADA, this page may be clipped and copied as a handout for patients, without first obtaining reprint permission from the ADA Publishing Division. Any other use, copying or distribution, whether in printed or electronic form, is strictly prohibited without prior written consent of the ADA Publishing Division.

"For the Dental Patient" provides general information on dental treatments to dental patients. It is designed to prompt discussion between dentist and patient about treatment options and does not substitute for the dentist's professional assessment based on the individual patient's needs and desires.